

LET'S TALK ABOUT HOW YOU CURRENTLY MANAGE MIGRAINE

It's time to tell your healthcare provider about what's working or not with your current migraine treatment plan.

Fill out the questions below and bring your answers to your next appointment.

How many migraine attacks do you have each month?



1



2



3



4



5



6



7+

What treatments—both prescription and over the counter—do you currently take for your migraine attacks when they strike?

Do you ever take multiple types of medication to treat a migraine attack?

Yes No

ASK YOUR HEALTHCARE PROVIDER: *Can I take UBRELVY with other migraine medications?*

Which symptoms do you experience during a migraine attack?



Head Pain



Light Sensitivity



Sound Sensitivity



Nausea



All of these

ASK YOUR HEALTHCARE PROVIDER: *Does UBRELVY treat migraine pain and symptoms?*

How often do you question taking your migraine medication because you're in a place where you feel you can't take it?

Never

For a few of my migraine attacks

For many of my migraine attacks

For all of my migraine attacks

ASK YOUR HEALTHCARE PROVIDER: *Are there any side effects with UBRELVY?*

Do you ever wait to see if your migraine attack will be "bad enough" before treating it with prescription medication? If so, do you still get the relief you need?

Never

For a few of my migraine attacks

For many of my migraine attacks

For all of my migraine attacks

ASK YOUR HEALTHCARE PROVIDER: *If I don't take my medication right away, will UBRELVY still work?*

After answering these questions, write down what is and isn't working with your current migraine management plan and any other questions you have about UBRELVY.

 **TAKE THIS WITH YOU AND ASK YOUR HEALTHCARE PROVIDER ABOUT UBRELVY TODAY!**

What is UBRELVY™ (ubrogepant)?

UBRELVY is a prescription medicine used for the acute treatment of migraine attacks with or without aura in adults. UBRELVY is not used to prevent migraine headaches.

IMPORTANT SAFETY INFORMATION

Who should not take UBRELVY™ (ubrogepant)?

Do not take UBRELVY if you are taking medicines known as strong CYP3A4 inhibitors, such as ketoconazole, clarithromycin, itraconazole.

Please see full Patient Information on the next page.

UBR131444-v2 06/20

Patient Information
UBRELVY™ (you-brel-vee)
(ubrogepant)
tablets, for oral use

What is UBRELVY?

UBRELVY is a prescription medicine used for the acute treatment of migraine attacks with or without aura in adults. UBRELVY is not used to prevent migraine headaches. It is not known if UBRELVY is safe and effective in children.

Do not take UBRELVY if you are taking medicines known as a strong CYP3A4 inhibitor, such as:

- ketoconazole
- clarithromycin
- itraconazole

Ask your healthcare provider if you are not sure if you are taking any of these medicines. Your healthcare provider can tell you if it is safe to take UBRELVY with other medicines.

Before you take UBRELVY tell your healthcare provider about all of your medical conditions, including if you:

- have liver problems
- have kidney problems
- are pregnant or plan to become pregnant. It is not known if UBRELVY will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if UBRELVY passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Especially tell your healthcare provider if you take any of the following, as your healthcare provider may need to change the dose of UBRELVY:

- verapamil
- cyclosporine
- ciprofloxacin
- fluconazole
- fluvoxamine
- phenytoin
- barbiturates
- rifampin
- St. John's Wort
- quinidine
- carvedilol
- eltrombopag
- curcumin

These are not all of the medicines that could affect how UBRELVY works. Your healthcare provider can tell you if it is safe to take UBRELVY with other medicines.

Keep a list of medicines you take to show to your healthcare provider or pharmacist when you get a new medicine.

How should I take UBRELVY?

- Take UBRELVY exactly as your healthcare provider tells you to take it.
- Take UBRELVY with or without food.
- Most patients can take a second tablet 2 hours after the first tablet, as needed.
- You should not take a second tablet within 24 hours if you consume grapefruit or grapefruit juice or are taking medications that may include:
 - verapamil
 - cyclosporin
 - ciprofloxacin
 - fluconazole
 - fluvoxamine
- It is not known if it is safe to take UBRELVY for more than 8 migraine headaches in 30 days.
- You should write down when you have headaches and when you take UBRELVY so you can talk to your healthcare provider about how UBRELVY is working for you.
- If you take too much UBRELVY, call your healthcare provider or go to the nearest hospital emergency room right away.

What are the possible side effects of UBRELVY?

The most common side effects of UBRELVY are nausea and sleepiness.

These are not all of the possible side effects of UBRELVY. Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Distributed by: Allergan USA, Inc.

Madison, NJ 07940

© 2019 Allergan. All rights reserved.

UBRELVY™ is a trademark of Allergan Pharmaceuticals International Limited.

Allergan® and its design are trademarks of Allergan, Inc.

Patented. See www.allergan.com/patents

